

بِسْمِ تَعَالَى



شرکت مدیریت منابع آب ایران

بررسی آثار احتمالی پساب حاوی ویروس

COVID-19

بر منابع آب

دفتر مطالعات پایه منابع آب - گروه محیط زیست و کیفیت منابع آب

فروردین ۱۳۹۹

- ملاحظات مربوط به اثرات احتمالی ویروس COVID-19 بر منابع آب سطحی و زیرزمینی که از توصیه‌های سازمان جهانی بهداشت^۱ (WHO) برگرفته شده، به شرح زیر ارائه می‌گردد. (توصیه‌های یاد شده پیوست می‌باشد)
- ۱- ویروس COVID-19، یک ویروس پوشش‌دار باغشای بیرونی شکننده است. بطور کلی ویروس‌های پوشش‌دار در محیط زیست از پایداری کمتری برخوردار بوده و بیشتر به اکسیدان‌هایی نظیر کلر حساس هستند.
 - ۲- در حالیکه هیچ مدرکی در مورد بقای ویروس COVID-19 در آب و فاضلاب وجود ندارد، لیکن احتمالاً این ویروس، بطور قابل توجهی سریع‌تر از ویروس‌های روده‌ای انسانی بدون پوشش که ناقل بیماری‌های شناخته شده ناشی از آب می‌باشند (نظیر آدوموویروس، نورو ویروس و هپاتیت A)، غیر فعال می‌شود.
 - ۳- تاکنون شواهد متقنی مبنی بر انتقال ویروس COVID-19 از طریق سیستم فاضلاب با یا بدون تصفیه و همچنین دفع در چاه‌های جاذب گزارش نشده است.
 - ۴- گرچه در برخی کشورها، اموات شستشو داده نمی‌شوند، اما وجود ویروس COVID-19 در پساب حاصله از تصفیل محتمل است.
 - ۵- برخی مطالعات نشان می‌دهد که یک کرونا ویروس انسانی تا ۲ روز در آب لوله کشی فاقد کلر و در فاضلاب بیمارستانی با دمای ۲۰ درجه سانتی گراد زنده مانده است. همچنین به استناد مقاله منتشره در مجله **Water research**^۲ ماندگاری گونه مشابه این ویروس در آب خام چندین روز و در فاضلاب چندین هفته اعلام شده است.
 - ۶- بنابراین بنظر می‌رسد برای نابودی ویروس COVID-19، استفاد از آب با غلظت کلر بالاتر از استاندارد آب آشامیدنی نیاز است که این موضوع می‌بایستی برای تعیین غلظت مناسب مورد بررسی و تحقیق قرار گیرد.

✚ برخی از مهمترین پیشنهادات دو مرجع فوق به منظور تعدیل اثرات ویروس بر منابع آب:

- ۱- گورهای محل تدفین اموات حداقل ۳۰ متر فاصله افقی از منابع آب زیرزمینی که برای آب آشامیدنی استفاده می‌شود، داشته باشند.
- ۲- کف گور، باید حداقل ۱۵۰ سانتیمتر بالاتر از تراز سطح آب زیرزمینی و حداقل عمق گور از ۷۰ سانتیمتر کمتر نباشد. (به عبارتی سطح آب زیرزمینی در محل گورستان از عمق حداقل ۲۲۰ سانتیمتر کمتر نباشد)

1 _Risks posed by dead bodies after disasters, Extract from WER No. 3, 2005, 80, 21-28, published by WHO Unit on Disease Control in Humanitarian Emergencies (https://www.who.int/diseasecontrol_emergencies/guidelines/risks/en/)

_Water, sanitation, hygiene and waste management for the COVID-19 virus, Technical brief, 3 March 2020

2.Casanova,L. , Rutala. W.A., Weber,D.J. and Sobsey,M.D., 2009.Survial of surrogate coronaviruses in water. Water research. 43(7), pp.1893-1898

۳- فاضلاب ناشی از تغسیل قبل از تصفیه و گندزدایی نباید وارد محیط های انسانی شود.

توصیه ها:

- ۱- آب مصرفی غسلخانه می بایستی از غلظت کلر بالاتر از استاندارد آب شرب برخوردار باشد.
- ۲- در قسمت تغسیل اموات، محلول ۰/۵ میلیگرم در لیتر آب ژاول بر روی جسد اسپری شود تا اطمینان حاصل گردد که ویروس از بین رفته است. این کار در درجه‌ی اول برای پرسنل غسلخانه ضروری است و در درجه دوم باعث می‌شود تا احتمال ورود ویروس به محیط زیست به حداقل کاهش یابد.
- ۳- از هرگونه استفاده از آب خام فاقد سیستم گندزدایی در غسلخانه ها خودداری گردد.
- ۴- از دفن اموات آلوده به کرونا در گورستانهایی که سطح برخورد آب زیرزمینی آنها کمتر از ۲۲۰ سانتی متر است، اکیدا جلوگیری گردد.
- ۵- فاضلاب ناشی از تغسیل قبل از تصفیه و گندزدایی نباید وارد محیط های انسانی شود و از هر گونه اقدام مغایر باید جلوگیری شود.
- ۶- در دفن اموات آلوده به کرونا، رعایت پروتکل های بهداشتی مربوطه لازم الرعایه است.

❖ با توجه به موارد نامبرده، بنظر میرسد غسل اموات آلوده به ویروس COVID-19 با رعایت نکات برشمردده فوق و نیز رعایت پروتکل های بهداشتی مربوط به دفن اموات عفونی، آثار نامطلوبی بر منابع آب وارد نکند.

پیشنهادات پایانی:

- ۱- نظر به اینکه وفق ماده ۴۶ قانون توزیع عادلانه آب، مسئولیت پیگیری، کنترل و ممانعت از آلودگی منابع آب با سازمان محیط زیست می‌باشد. لذا ضروری است تا این مهم از سوی سازمان مذکور با هماهنگی وزارت بهداشت و درمان و آموزش پزشکی مورد پیگیری و اقدام قرار گیرد.
- ۲- رفع آلودگی ناشی از سیستم دفع فاضلاب بیمارستانی و بویژه بیمارستان‌های فاقد سیستم تصفیه فاضلاب، باید در اولویت قرار گیرد.
- ۳- برای بررسی همه جانبه اثرات این ویروس بر منابع آب، در قالب طرح تحقیقات کاربردی این مهم جهت بررسی بیشتر به دانشکده بهداشت محیط دانشگاه تهران، شهید بهشتی و ... ارجاع گردد.

Water, sanitation, hygiene, and waste management for the COVID-19 virus

Interim guidance
19 March 2020

Background

This interim guidance supplements the infection prevention and control (IPC) documents by summarizing WHO guidance on water, sanitation and health care waste relevant to viruses, including coronaviruses. It is intended for water and sanitation practitioners and providers and health care providers who want to know more about water, sanitation and hygiene (WASH) risks and practices.

The provision of safe water, sanitation, and hygienic conditions is essential to protecting human health during all infectious disease outbreaks, including the COVID-19 outbreak. Ensuring good and consistently applied WASH and waste management practices in communities, homes, schools, marketplaces, and health care facilities will help prevent human-to-human transmission of the COVID-19 virus.

The most important information concerning WASH and the COVID-19 virus is summarized here.

- Frequent and proper hand hygiene is one of the most important measures that can be used to prevent infection with the COVID-19 virus. WASH practitioners should work to enable more frequent and regular hand hygiene by improving facilities and using proven behavior-change techniques.
- WHO guidance on the safe management of drinking-water and sanitation services applies to the COVID-19 outbreak. Extra measures are not needed. Disinfection will facilitate more rapid die-off of the COVID-19 virus.
- Many co-benefits will be realized by safely managing water and sanitation services and applying good hygiene practices.

Currently, there is no evidence about the survival of the COVID-19 virus in drinking-water or sewage. The morphology and chemical structure of the COVID-19 virus are similar to those of other human coronaviruses for which there are data about both survival in the environment and effective inactivation measures. This document draws upon the evidence base and WHO guidance on how to protect against viruses in sewage and drinking-water. This document will be updated as new information becomes available.

1. COVID-19 transmission

There are two main routes of transmission of the COVID-19 virus: respiratory and contact. Respiratory droplets are generated when an infected person coughs or sneezes. Any person who is in close contact with someone who has respiratory symptoms (sneezing, coughing) is at risk of being exposed to potentially infective respiratory droplets.¹ Droplets may also land on surfaces where the virus could remain viable; thus, the immediate environment of an infected individual can serve as a source of transmission (contact transmission).

Approximately 2–10% of cases of confirmed COVID-19 disease present with diarrhoea,^{2,4} and two studies detected COVID-19 viral RNA fragments in the faecal matter of COVID-19 patients.^{5,6} However, only one study has cultured the COVID-19 virus from a single stool specimen.⁷ There have been no reports of faecal–oral transmission of the COVID-19 virus.

2. Persistence of the COVID-19 virus in drinking-water, faeces and sewage and on surfaces.

Although persistence in drinking-water is possible, there is no evidence from surrogate human coronaviruses that they are present in surface or groundwater sources or transmitted through contaminated drinking water. The COVID-19 virus is an enveloped virus, with a fragile outer membrane. Generally, enveloped viruses are less stable in the environment and are more susceptible to oxidants, such as chlorine. While there is no evidence to date about survival of the COVID-19 virus in water or sewage, the virus is likely to become inactivated significantly faster than non-enveloped human enteric viruses with known waterborne transmission (such as adenoviruses, norovirus, rotavirus and hepatitis A). For example, one study found that a surrogate human coronavirus survived only 2 days in dechlorinated tap water and in hospital wastewater at 20°C.⁸ Other studies concur, noting that the human coronaviruses transmissible gastroenteritis coronavirus and mouse hepatitis virus demonstrated a 99.9% die-off in from 2 days⁹ at 23°C to 2 weeks¹⁰ at 25°C. Heat, high or low pH, sunlight, and common disinfectants (such as chlorine) all facilitate die off.

It is not certain how long the virus that causes COVID-19 survives on surfaces, but it seems likely to behave like other coronaviruses. A recent review of the survival of human

coronaviruses on surfaces found large variability, ranging from 2 hours to 9 days.¹¹ The survival time depends on a number of factors, including the type of surface, temperature, relative humidity, and specific strain of the virus. The same review also found that effective inactivation could be achieved within 1 minute using common disinfectants, such as 70% ethanol or sodium hypochlorite (for details, see Cleaning practices).

3. Keeping water supplies safe

The COVID-19 virus has not been detected in drinking-water supplies, and based on current evidence, the risk to water supplies is low.¹² Laboratory studies of surrogate coronaviruses that took place in well-controlled environments indicated that the virus could remain infectious in water contaminated with faeces for days to weeks.¹⁰ A number of measures can be taken to improve water safety, starting with protecting the source water; treating water at the point of distribution, collection, or consumption; and ensuring that treated water is safely stored at home in regularly cleaned and covered containers.

Conventional, centralized water treatment methods that use filtration and disinfection should inactivate the COVID-19 virus. Other human coronaviruses have been shown to be sensitive to chlorination and disinfection with ultraviolet (UV) light.¹³ As enveloped viruses are surrounded by a lipid host cell membrane, which is not robust, the COVID-19 virus is likely to be more sensitive to chlorine and other oxidant disinfection processes than many other viruses, such as coxsackieviruses, which have a protein coat. For effective centralized disinfection, there should be a residual concentration of free chlorine of ≥ 0.5 mg/L after at least 30 minutes of contact time at pH < 8.0 .¹² A chlorine residual should be maintained throughout the distribution system.

In places where centralized water treatment and safe piped water supplies are not available, a number of household water treatment technologies are effective in removing or destroying viruses, including boiling or using high-performing ultrafiltration or nanomembrane filters, solar irradiation and, in non-turbid waters, UV irradiation and appropriately dosed free chlorine.

4. Safely managing wastewater and faecal waste

There is no evidence that the COVID-19 virus has been transmitted via sewerage systems with or without wastewater treatment. Further, there is no evidence that sewage or wastewater treatment workers contracted the severe acute respiratory syndrome (SARS), which is caused by another type of coronavirus that caused a large outbreak of acute respiratory illness in 2003. As part of an integrated public health policy, wastewater carried in sewerage systems should be treated in well-designed and well-managed centralized wastewater treatment works. Each stage of treatment (as well as retention time and dilution) results in a further reduction of the potential risk. A waste stabilization pond (an oxidation pond or lagoon) is generally considered a practical and simple wastewater treatment technology particularly well suited to destroying pathogens, as relatively long retention times (20 days or longer) combined with sunlight, elevated pH levels, biological activity, and other factors serve to accelerate pathogen destruction. A final disinfection step may be considered if existing wastewater treatment plants are not optimized to remove viruses. Best practices for protecting the health of workers at sanitation treatment facilities should

be followed. Workers should wear appropriate personal protective equipment (PPE), which includes protective outerwear, gloves, boots, goggles or a face shield, and a mask; they should perform hand hygiene frequently; and they should avoid touching eyes, nose, and mouth with unwashed hands.

WASH in health care settings

Existing recommendations for water, sanitation and hygiene measures in health care settings are important for providing adequate care for patients and protecting patients, staff, and caregivers from infection risks.¹⁴ The following actions are particularly important: (i) managing excreta (faeces and urine) safely, including ensuring that no one comes into contact with it and that it is treated and disposed of correctly; (ii) engaging in frequent hand hygiene using appropriate techniques; (iii) implementing regular cleaning and disinfection practices; and (iv) safely managing health care waste. Other important measures include providing sufficient safe drinking-water to staff, caregivers, and patients; ensuring that personal hygiene can be maintained, including hand hygiene, for patients, staff and caregivers; regularly laundering bedsheets and patients' clothing; providing adequate and accessible toilets (including separate facilities for confirmed and suspected cases of COVID-19 infection); and segregating and safely disposing of health care waste. For details on these recommendations, please refer to Essential environmental health standards in health care.¹⁴

1. Hand hygiene practices

Hand hygiene is extremely important. Cleaning hands with soap and water or an alcohol-based hand rub should be performed according to the instructions known as “My 5 moments for hand hygiene”.¹⁵ If hands are not visibly dirty, the preferred method is to perform hand hygiene with an alcohol-based hand rub for 20–30 seconds using the appropriate technique.¹⁶ When hands are visibly dirty, they should be washed with soap and water for 40–60 seconds using the appropriate technique.¹⁷ Hand hygiene should be performed at all five moments, including before putting on PPE and after removing it, when changing gloves, after any contact with a patient with suspected or confirmed COVID-19 infection or their waste, after contact with any respiratory secretions, before eating, and after using the toilet.¹⁸ If an alcohol-based hand rub and soap are not available, then using chlorinated water (0.05%) for handwashing is an option, but it is not ideal because frequent use may lead to dermatitis, which could increase the risk of infection and asthma and because prepared dilutions might be inaccurate.¹⁹ However, if other options are not available or feasible, using chlorinated water for handwashing is an option.

Functional hand hygiene facilities should be present for all health care workers at all points of care and in areas where PPE is put on or taken off. In addition, functional hand hygiene facilities should be available for all patients, family members, and visitors, and should be available within 5 m of toilets, as well as in waiting and dining rooms and other public areas.

2. Sanitation and plumbing

People with suspected or confirmed COVID-19 disease should be provided with their own flush toilet or latrine that has a door that closes to separate it from the patient's room. Flush toilets should operate properly and have functioning drain traps. When possible, the toilet should be flushed with the lid down to prevent droplet splatter and aerosol clouds. If it is not possible to provide separate toilets, the toilet should be cleaned and disinfected at least twice daily by a trained cleaner wearing PPE (gown, gloves, boots, mask, and a face shield or goggles). Further, and consistent with existing guidance, staff and health care workers should have toilet facilities that are separate from those used by all patients.

WHO recommends the use of standard, well-maintained plumbing, such as sealed bathroom drains, and backflow valves on sprayers and faucets to prevent aerosolized faecal matter from entering the plumbing or ventilation system,²⁰ together with standard wastewater treatment.²¹ Faulty plumbing and a poorly designed air ventilation system were implicated as contributing factors to the spread of the aerosolized SARS coronavirus in a high-rise apartment building in Hong Kong in 2003.²² Similar concerns have been raised about the spread of the COVID-19 virus from faulty toilets in high-rise apartment buildings.²³ If health care facilities are connected to sewers, a risk assessment should be conducted to confirm that wastewater is contained within the system (that is, the system does not leak) before its arrival at a functioning treatment or disposal site, or both. Risks pertaining to the adequacy of the collection system or to treatment and disposal methods should be assessed following a safety planning approach,²⁴ with critical control points prioritized for mitigation.

For smaller health care facilities in low-resource settings, if space and local conditions allow, pit latrines may be the preferred option. Standard precautions should be taken to prevent contamination of the environment by excreta. These precautions include ensuring that at least 1.5 m exists between the bottom of the pit and the groundwater table (more space should be allowed in coarse sands, gravels, and fissured formations) and that the latrines are located at least 30 m horizontally from any groundwater source (including both shallow wells and boreholes).²¹ If there is a high groundwater table or a lack of space to dig pits, excreta should be retained in impermeable storage containers and left for as long as feasible to allow for a reduction in virus levels before moving it off-site for additional treatment or safe disposal, or both. A two-tank system with parallel tanks would help facilitate inactivation by maximizing retention times, as one tank could be used until full, then allowed to sit while the next tank is being filled. Particular care should be taken to avoid splashing and the release of droplets while cleaning or emptying tanks.

3. Toilets and the handling of faeces

It is critical to conduct hand hygiene when there is suspected or direct contact with faeces (if hands are dirty, then soap and water are preferred to the use of an alcohol-based hand rub). If the patient is unable to use a latrine, excreta should be collected in either a diaper or a clean bedpan and immediately and carefully disposed of into a separate toilet or latrine used only by suspected or confirmed cases of COVID-19. In all health care settings, including those with suspected or confirmed COVID-19 cases, faeces must be treated as a biohazard and handled as little as possible. Anyone handling

faeces should follow WHO contact and droplet precautions¹⁸ and use PPE to prevent exposure, including long-sleeved gowns, gloves, boots, masks, and goggles or a face shield. If diapers are used, they should be disposed of as infectious waste as they would be in all situations. Workers should be properly trained in how to put on, use, and remove PPE so that these protective barriers are not breached.²⁵ If PPE is not available or the supply is limited, hand hygiene should be regularly practiced, and workers should keep at least 1 m distance from any suspected or confirmed cases.

If a bedpan is used, after disposing of excreta from it, the bedpan should be cleaned with a neutral detergent and water, disinfected with a 0.5% chlorine solution, and then rinsed with clean water; the rinse water should be disposed of in a drain or a toilet or latrine. Other effective disinfectants include commercially available quaternary ammonium compounds, such as cetylpyridinium chloride, used according to manufacturer's instructions, and peracetic or peroxyacetic acid at concentrations of 500–2000 mg/L.²⁶

Chlorine is ineffective for disinfecting media containing large amounts of solid and dissolved organic matter. Therefore, there is limited benefit to adding chlorine solution to fresh excreta and it is possible that this may introduce risks associated with splashing.

4. Emptying latrines and holding tanks, and transporting excreta off-site.

There is no reason to empty latrines and holding tanks of excreta from suspected or confirmed COVID-19 cases unless they are at capacity. In general, the best practices for safely managing excreta should be followed. Latrines or holding tanks should be designed to meet patient demand, considering potential sudden increases in cases, and there should be a regular schedule for emptying them based on the wastewater volumes generated. PPE (long-sleeved gown, gloves, boots, masks, and goggles or a face shield) should be worn at all times when handling or transporting excreta offsite, and great care should be taken to avoid splashing. For crews, this includes pumping out tanks or unloading pumper trucks. After handling the waste and once there is no risk of further exposure, individuals should safely remove their PPE and perform hand hygiene before entering the transport vehicle. Soiled PPE should be put in a sealed bag for later safe laundering (see Cleaning practices). Where there is no off-site treatment, in-situ treatment can be done using lime. Such treatment involves using a 10% lime slurry added at 1-part lime slurry per 10 parts of waste.

5. Cleaning practices

Recommended cleaning and disinfection procedures for health care facilities should be followed consistently and correctly.¹⁹ Laundry should be done and surfaces in all environments in which COVID-19 patients receive care (treatment units, community care centres) should be cleaned at least once a day and when a patient is discharged.²⁷ Many disinfectants are active against enveloped viruses, such as the COVID-19 virus, including commonly used hospital disinfectants. Currently, WHO recommends using:

- 70% ethyl alcohol to disinfect small areas between uses, such as reusable dedicated equipment (for example, thermometers);
- sodium hypochlorite at 0.5% (equivalent to 5000 ppm) for disinfecting surfaces.

All individuals dealing with soiled bedding, towels, and clothes from patients with COVID-19 infection should wear appropriate PPE before touching soiled items, including heavy duty gloves, a mask, eye protection (goggles or a face shield), a long-sleeved gown, an apron if the gown is not fluid resistant, and boots or closed shoes. They should perform hand hygiene after exposure to blood or body fluids and after removing PPE. Soiled linen should be placed in clearly labelled, leak-proof bags or containers, after carefully removing any solid excrement and putting it in a covered bucket to be disposed of in a toilet or latrine. Machine washing with warm water at 60–90°C (140–194°F) with laundry detergent is recommended. The laundry can then be dried according to routine procedures. If machine washing is not possible, linens can be soaked in hot water and soap in a large drum using a stick to stir and being careful to avoid splashing. The drum should then be emptied, and the linens soaked in 0.05% chlorine for approximately 30 minutes. Finally, the laundry should be rinsed with clean water and the linens allowed to dry fully in sunlight.

If excreta are on surfaces (such as linens or the floor), the excreta should be carefully removed with towels and immediately safely disposed of in a toilet or latrine. If the towels are single use, they should be treated as infectious waste; if they are reusable, they should be treated as soiled linens. The area should then be cleaned and disinfected (with, for example, 0.5% free chlorine solution), following published guidance on cleaning and disinfection procedures for spilled body fluids.²⁷

6. Safely disposing of greywater or water from washing PPE, surfaces and floors.

Current WHO recommendations are to clean utility gloves or heavy duty, reusable plastic aprons with soap and water and then decontaminate them with 0.5% sodium hypochlorite solution after each use. Single-use gloves (nitrile or latex) and gowns should be discarded after each use and not reused; hand hygiene should be performed after PPE is removed. If greywater includes disinfectant used in prior cleaning, it does not need to be chlorinated or treated again. However, it is important that such water is disposed of in drains connected to a septic system or sewer or in a soakaway pit. If greywater is disposed of in a soakaway pit, the pit should be fenced off within the health facility grounds to prevent tampering and to avoid possible exposure in the case of overflow.

7. Safe management of health care waste

Best practices for safely managing health care waste should be followed, including assigning responsibility and sufficient human and material resources to dispose of such waste safely. There is no evidence that direct, unprotected human contact during the handling of health care waste has resulted in the transmission of the COVID-19 virus. All health care waste produced during the care of COVID 19 patients should be collected safely in designated containers and bags, treated, and then safely disposed of or treated, or both, preferably on-site. If waste is moved off-site, it is critical to understand where and how it will be treated and destroyed. All who handle health care waste should wear appropriate PPE (boots, apron, long-sleeved gown, thick gloves, mask, and goggles or a face shield) and perform hand hygiene after removing it. For more information refer to the WHO guidance, Safe management of wastes from health-care activities.²⁸

Considerations for WASH practices in homes and communities.

Upholding best WASH practices in the home and community is also important for preventing the spread of COVID-19 and when caring for patients at home. Regular and correct hand hygiene is of particular importance.

1. Hand hygiene

Hand hygiene in non-health care settings is one of the most important measures that can prevent COVID 19 infection. In homes, schools and crowded public spaces – such as markets, places of worship, and train or bus stations – regular handwashing should occur before preparing food, before and after eating, after using the toilet or changing a child’s diaper, and after touching animals. Functioning handwashing facilities with water and soap should be available within 5 m of toilets.

2. Treatment and handling requirements for excreta.

Best WASH practices, particularly handwashing with soap and clean water, should be strictly applied and maintained because these provide an important additional barrier to COVID-19 transmission and to the transmission of infectious diseases in general.¹⁷ Consideration should be given to safely managing human excreta throughout the entire sanitation chain, starting with ensuring access to regularly cleaned, accessible, and functioning toilets or latrines and to the safe containment, conveyance, treatment, and eventual disposal of sewage.

When there are suspected or confirmed cases of COVID-19 in the home setting, immediate action must be taken to protect caregivers and other family members from the risk of contact with respiratory secretions and excreta that may contain the COVID-19 virus. Frequently touched surfaces throughout the patient’s care area should be cleaned regularly, such as beside tables, bed frames and other bedroom furniture. Bathrooms should be cleaned and disinfected at least once a day. Regular household soap or detergent should be used for cleaning first and then, after rinsing, regular household disinfectant containing 0.5% sodium hypochlorite (that is, equivalent to 5000 ppm or 1-part household bleach with 5% sodium hypochlorite to 9 parts water) should be applied. PPE should be worn while cleaning, including mask, goggles, a fluid-resistant apron, and gloves,²⁹ and hand hygiene with an alcohol-based hand rub or soap and water should be performed after removing PPE.

References

1. Coronavirus disease (COVID-19) advice for the public. Geneva: World Health Organization; 2020 (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>, accessed 3 March 2020).
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506. doi:10.1016/S0140-6736(20)30183-5.

3. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395:507–13. doi:10.1016/S0140-6736(20)30211-7.
4. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020. Feb 7. doi:10.1001/jama.2020.1585.
5. Xiao E, Tang M, Zheng Y, Li C, He J, Hong H, et al. Evidence for gastrointestinal infection of SARS-CoV. medRxiv. doi:10.1101/2020.02.17.20023721.
6. Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H et al. for the Washington State 2019-nCoV Case Investigation Team. First case of 2019 novel coronavirus in the United States. *N Engl J Med*. 2020. Jan 31. doi:10.1056/NEJMoa2001191.
7. Zhang Y, Chen C, Zhu S et al. [Isolation of 2019-nCoV from a stool specimen of a laboratory-confirmed case of the coronavirus disease 2019 (COVID-19)]. *China CDC Weekly*. 2020;2(8):123-4. (In Chinese.)
8. Wang XW, Li JS, Zhen B, Kong QX, Song N, Xiao WJ et al. Study on the resistance of severe acute respiratory syndrome-associated coronavirus. *J Virol Methods*. 2005;126:171–7. doi:10.1016/j.jviromet.2005.02.005.
9. Gundy P, Gerba CP, Pepper IL. Survival of coronaviruses in water and wastewater. *Food Environ Virol*. 2009;1:10-14. doi:10.1007/s12560-008-9001-6.
10. Casanova L, Rutalal WA, Weber DJ, Sobsey MD. Survival of surrogate coronaviruses in water. *Water Res*. 2009;43(7):1893–8. doi:10.1016/j.watres.2009.02.002.
11. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect*. 2020;104(3):246–51. doi:10.1016/j.jhin.2020.01.022.
12. Guidelines for drinking-water quality, fourth edition, incorporating the first addendum. Geneva: World Health Organization; 2017 (<http://apps.who.int/iris/bitstream/10665/254637/1/9789241549950-eng.pdf>, accessed 3 March 2020).
13. SARS-CoV-2 – water and sanitation. Adelaide: Water Research Australia; 2020 (http://www.waterra.com.au/r9544/media/system/attrib/file/2199/WaterRA_FS_Coronavirus_V10.pdf, accessed 3 March 2020).
14. Essential environmental health standards in health care. Geneva: World Health Organization; 2008 (https://apps.who.int/iris/bitstream/handle/10665/43767/9789241547239_eng.pdf?sequence=1&isAllowed=y, accessed 3 March 2020).
15. My 5 moments for hand hygiene. In: WHO/Infection prevention and control [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/infection-prevention/campaigns/clean-hands/5moments/en/>, accessed 3 March 2020).
16. Siddharta A, Pfaender S, Vielle NJ, Dijkman R, Friesland M, Becker B, et al. Virucidal activity of World Health Organization-recommended formulations against enveloped viruses, including Zika, Ebola, and emerging coronaviruses. *J Infect Dis*. 2017;215(6):902–6. doi:10.1093/infdis/jix046.
17. WHO guidelines on hand hygiene in health care settings. Geneva: World Health Organization; 2009 (https://apps.who.int/iris/bitstream/handle/10665/44102/9789241597906_eng.pdf?sequence=1&isAllowed=y, accessed 3 March 2020).
18. Infection prevention and control during health care when novel coronavirus (nCoV) infection is suspected: interim guidance, 25 January 2020. Geneva: World Health Organization ([https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-\(ncov\)-infection-is-suspected-20200125](https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-(ncov)-infection-is-suspected-20200125), accessed 3 March 2020).
19. Q&A on infection prevention and control for health care workers caring for patients with suspected or confirmed 2019-nCoV. In: WHO/Newsroom [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/news-room/q-a-detail/q-a-on-infection-prevention-and-control-for-health-care-workers-caring-for-patients-with-suspected-or-confirmed-2019-ncov>, accessed 3 March 2020).
20. Health aspects of plumbing. Geneva: World Health Organization; 2006. (<https://apps.who.int/iris/handle/10665/43423>, accessed 3 March 2020).
21. Guidelines on sanitation and health. Geneva: World Health Organization; 2018 (<https://apps.who.int/iris/bitstream/handle/10665/274939/9789241514705-eng.pdf?ua=1>, accessed 3 March 2020).
22. Yu ITS, Li Y, Wong TW, Tam W, Chan A, Lee JHW, et al. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N Engl J Med*. 2004;350(17): 1731-9. doi:10.1056/NEJMoa032867.
23. Regan H. How can the coronavirus spread through bathroom pipes? Experts are investigating in Hong Kong. CNN. 12 February 2020 (<https://edition.cnn.com/2020/02/12/asia/hong-kong-coronavirus-pipes-intl-hnk/index.html>).
24. Sanitation safety planning: manual for safe use and disposal of wastewater, greywater and excreta. Geneva: World Health Organization; 2015. (<https://apps.who.int/iris/handle/10665/171753>, accessed 3 March 2020).
25. How to put on and take off personal protective equipment. Geneva: World Health Organization; 2008 (<https://apps.who.int/iris/handle/10665/70066>, accessed 3 March 2020).
26. Chemical disinfectants: guideline for disinfection and sterilization in healthcare facilities (2008). In: CDC/Infection Control [website]. Atlanta: US Centers for Disease Control and Prevention; 2019. (<https://www.cdc.gov/infectioncontrol/guidelines/disinfection/disinfection-methods/chemical.html>, accessed 3 March 2020).

27. Best practices for environmental cleaning in healthcare facilities in resource-limited settings. Atlanta: US Centers for Disease Control and Prevention; 2019
(<https://www.cdc.gov/hai/pdfs/resource-limited/environmental-cleaning-508.pdf>, accessed 3 March 2020).
28. Safe management of wastes from health-care activities: a summary. Geneva: World Health Organization; 2017
(<https://apps.who.int/iris/handle/10665/259491>, accessed 3 March 2020).
29. Home care for patients with suspected novel coronavirus (COVID-19) infection presenting with mild symptoms, and management of their contacts: interim guidance, 4 February 2020.
([https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-\(ncov\)-infection-presenting-with-mild-symptoms-and-management-of-contacts](https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-(ncov)-infection-presenting-with-mild-symptoms-and-management-of-contacts), accessed 3 March 2020).

US Centers for Disease Control and Prevention, United States of America; David Berendes, US Centers for Disease Control and Prevention, United States of America; Lisa Casanova, Georgia State University, United States of America; David Cunliffe, SA Health, Australia; Rick Gelting, US Centers for Disease Control and Prevention, United States of America; Dr Thomas Handzel, US Centers for Disease Control and Prevention, United States of America; Paul Hunter, University of East Anglia, United Kingdom; Ana Maria de Roda Husman, National Institute for Public Health and the Environment, the Netherlands; Peter Maes, Médecins Sans Frontières, Belgium; Molly Patrick, US Centers for Disease Control and Prevention, United States of America; Mark Sobsey, University of North Carolina-Chapel Hill, United States of America.

WHO continues to monitor the situation closely for any changes that may affect this interim guidance. Should any factors change, WHO will issue a further update. Otherwise, this interim guidance document will expire 2 years after the date of publication.

Contributors

This interim guidance was written by staff from WHO and UNICEF. In addition, a number of experts and WASH practitioners contributed. They include Matt Arduino,

© World Health Organization 2020. Some rights reserved. This work is available under the [CC BY-NC-SA 3.0 IGO](https://creativecommons.org/licenses/by-nc-sa/3.0/) licence.

Risks posed by dead bodies after disasters

Contrary to common belief, there is no evidence that corpses pose a risk of epidemic disease after a natural disaster. Most agents do not survive long in the human body after death. Human remains only pose a substantial risk to health in a few special cases, such as deaths from cholera or haemorrhagic fevers.

Workers who routinely **handle** corpses may however risk contracting tuberculosis, bloodborne viruses (eg hepatitis B and C and HIV) and gastrointestinal infections (e.g. cholera, *E. coli*, hepatitis A, rotavirus diarrhoea, salmonellosis, shigellosis and typhoid/paratyphoid fevers):

- Tuberculosis can be acquired if the bacillus is aerosolized – residual air in lungs exhaled, fluid from lungs spurted up through the nose or mouth during handling of the corpse.
- Bloodborne viruses can be transmitted via direct contact of non-intact skin or mucous membrane from splashing of blood or body fluid or from injury from bone fragments and needles.
- Gastrointestinal (GI) infections can easily be transmitted from faeces leaked from dead bodies. Transmission occurs via the faecal–oral route through direct contact with the body, soiled clothes or contaminated vehicles or equipment. GI infections can also be spread as a result of contamination of the water supply with dead bodies.

Information on these risks should be provided to both emergency workers and the general public to ensure adequate disposal of bodies, appropriate precautions when handling bodies and to avoid panic and misunderstanding.

General advice

- In the case of mass casualties and where identification of victims is no longer possible, burial is preferable to cremation.
- Burial in mass graves is not a recommended public health measure. It can violate important social norms and waste scarce resources.
- The families' needs and social customs for funerals should be respected. If customs vary, each social group should be provided with a designated area, with the relevant materials, to be able to exercise their own traditions with dignity.
- Where existing facilities such as graveyards or crematoria are inadequate, alternative facilities should be provided.

Specific advice for workers handling corpses

- Graveyards should be at least 30 m from groundwater sources used for drinking-water.
- Grave floors must be at least 1.5 m above the water table, with a 0.7 m unsaturated zone.
- Surface water from graveyards must not enter inhabited areas.
- Exercise universal precautions taken when handling blood and body fluids.
- Use gloves once only and dispose of correctly.
- Use body bags.
- Wash hands with soap after handling bodies and before eating.
- Disinfect vehicles and equipment.
- Be vaccinated against hepatitis B.
- There is no need to disinfect bodies before disposal (except in case of cholera).

Risques liés aux cadavres après les désastres

Contrairement à une croyance répandue, rien ne prouve que les cadavres constituent un risque épidémique après une catastrophe naturelle. La plupart des agents infectieux ne survivent pas longtemps dans l'organisme humain après le décès. Les restes humains ne présentent un risque sanitaire que dans quelques cas particuliers nécessitant des précautions spéciales, comme les décès dus au choléra ou aux fièvres hémorragiques.

Toutefois, le personnel normalement appelé à **manipuler** des cadavres peut être exposé à la tuberculose, aux virus à transmission sanguine (hépatite B/C et VIH) et aux infections gastro-intestinales (choléra, *E. coli*, hépatite A, diarrhée à rotavirus, salmonellose, shigellose et fièvres typhoïde/paratyphoïde):

- Il est possible de contracter la tuberculose si le bacille est présent en aérosol (rejet de l'air résiduel des poumons, rejet par le nez ou la bouche de liquide provenant des poumons pendant la manipulation du cadavre).
- L'exposition à des virus à transmission sanguine résulte du contact direct entre une peau qui n'est pas intacte et du sang ou des liquides organiques, de traumatismes causés par des fragments osseux ou des aiguilles, ou de l'exposition des muqueuses à des éclaboussures de sang ou de liquides organiques.
- Les infections gastro-intestinales sont plus fréquentes car il n'est pas rare que des matières fécales émanent encore des cadavres. La transmission féco-orale résulte de contacts directs avec le cadavre, des vêtements souillés, ou encore des véhicules ou du matériel contaminés. Les cadavres qui contaminent les sources d'approvisionnement en eau peuvent aussi être à l'origine d'infections gastro-intestinales.

Il est important d'informer dûment le public et le personnel d'urgence afin d'éviter la panique et l'enterrement inapproprié des cadavres, et d'assurer que les cadavres sont manipulés avec les précautions voulues (voir la rubrique manipulation des cadavres ci-après).

Conseil général

- Lorsque les victimes sont nombreuses et que leur identification est impossible, l'enterrement est préférable à la crémation.
- Les corps ne doivent pas être placés dans des fosses communes sans cérémonie. Cette pratique viole des normes sociales importantes et peut représenter un gaspillage de ressources rares.
- Les familles devraient avoir la possibilité d'organiser des funérailles adaptées à leur culture et conformes à leurs coutumes. En présence de cultures différentes, chaque groupe social devrait disposer de zones distinctes et du matériel nécessaire pour observer ses traditions dignement.
- Faute d'installations satisfaisantes, qu'il s'agisse du cimetière ou du crématorium, d'autres sites et installations devront être trouvés.

Manipulation des cadavres

- Les cimetières doivent être à 30 mètres au moins des sources d'eau souterraines qui fournissent l'eau de boisson.
- Le fond d'une tombe doit être à 1,5 mètre au moins au-dessus de la nappe phréatique, avec une zone insaturée de 0,7 mètre.
- Les eaux superficielles provenant d'un cimetière ne doivent pas pénétrer dans les zones habitées.
- Observer les précautions universelles applicables au sang et aux liquides organiques.
- Utiliser des gants à usage unique (jetables) et à les éliminer correctement.
- Utiliser des sacs mortuaires.
- Se laver les mains avec du savon après avoir manipulé des cadavres et avant de manger.
- Désinfecter les véhicules et le matériel.
- Vacciner le personnel contre l'hépatite B.
- Il n'est pas utile de désinfecter les corps avant leur enterrement (sauf en cas de choléra).